

## The Cicada Parasite Beetles (Coleoptera: Rhipiceridae) of Virginia

Arthur V. Evans

1600 Nottoway Avenue  
Richmond, Virginia 23227  
arthurevans@verizon.net

Brent W. Steury

United States National Park Service  
700 George Washington Memorial Parkway  
Turkey Run Park Headquarters  
McLean, Virginia 22101  
Brent\_Steury@nps.gov

### ABSTRACT

The family Rhipiceridae is represented in Virginia by two species, *Sandalus niger* Knoch and *S. petrophya* Knoch, the latter of which is documented from the state for the first time.

*Key words:* cicada parasite beetle, new state record, Rhipiceridae, *Sandalus*, Virginia.

### INTRODUCTION

The Rhipiceridae, also known as cicada parasite or cedar beetles, is a small family with more than 100 species in seven genera (Lawrence, 2005). The family is represented in the Nearctic region by *Sandalus* (Greek σάνδαλον, or sandals), a genus proposed by Knoch (1801) to include two North American species *S. petrophya* and *S. niger*. *Sandalus* occurs in the New World, Africa, southeast Asia, China, India, and Japan (Katovich, 2002) and is in need of revision. The Nearctic fauna currently consists of five species, three of which occur in eastern North America. Of these, only two, *S. niger* Knoch and *S. petrophya* Knoch, occur in Virginia. The third species, *S. porosus* LeConte, ranges from Florida west to Colorado and Arizona (Katovich, 2002).

The common name “cedar beetle” was most likely instigated by LeConte’s (1862) statement that the Rhipiceridae was “A family containing a small number of species, found on plants; *Sandalus* especially affecting various cedars...” and popularized by Blatchley (1910) and Arnett (1963). The more recent moniker “cicada parasite beetles,” first coined by Downie & Arnett (1996), is more descriptive given the

available information on the larval biology of *Sandalus*. As noted by Dodge (1941), the “apparent association with certain trees has no significance except as it may indicate host or oviposition preferences of the host cicada.”

### METHODS

This study is based on our own field work, literature records, and the examination of specimens housed in the following collections: Virginia Museum of Natural History, Martinsville, Virginia (VMNH); Virginia Polytechnic Institute and State University, Blacksburg, Virginia (VPIC); National Museum of Natural History, Smithsonian Institution, Washington, D.C. (NMNH); George Washington Memorial Parkway, Turkey Run Park, McLean, Virginia (GWMP); Arthur V. Evans, Richmond, Virginia (AVEC).

### RESULTS AND DISCUSSION

Adult *Sandalus* are long (15-24 mm), convex, reddish brown or black (sometimes bicolored) and coarsely punctured beetles (Fig. 1); females are typically larger than males. The head has bulging eyes,



Fig. 1. *Sandalus petrophya* Knoch, female.

prominent hypognathous mandibles, and 11-segmented antennae that are distinctly flabellate in males and more or less serrate in females. The prothorax becomes wider posteriorly and is narrower than the base of the elytra. The long, vaguely ribbed and coarsely punctured elytra completely conceal the abdomen, which has five visible sternites. The tarsal formula is 5-5-5. Tarsomeres 1-4 are heart-shaped with membranous lobes and the claws are simple and equal in size.

Very little is known about the biology of rhipicerid beetles, with the exception of *S. niger*. Adults of this species do not feed and reach peak activity from late September through early October (Rings, 1942; Elzinga, 1977; Katovich 2002). They are found resting on tree trunks or grass, in Malaise trap samples, and are occasionally attracted to lights.

Elzinga (1977) observed mating aggregations of *S. niger* on the trunks of American Elm (*Ulmus americanus*), Shingle Oak (*Quercus imbricarius*), beech (*Fagus*), ash (*Fraxinus*), and other hardwoods. Adults emerge from their burrows in the morning and crawl up trunks to mate. Females lay large numbers of eggs in the holes and cracks of bark, preferably in areas where there are numerous cicadas. Rings (1942) noted that a single female could produce more than 16,000 eggs. Craighead (1921) proposed that the larvae of *S. niger* develop as ectoparasitoids of immature cicadas after discovering a pupa and cast larval exoskeleton of the beetle within a dead and hollowed-out cicada nymph. This assertion was later supported by descriptions of the first instar triungulin of *S. niger* (Dodge, 1941; Elzinga, 1977) that is typical of other beetle parasitoids. The larval stages between the triungulin and pupa remain unknown.

The adults of *Sandalus* in Virginia are distinguished by the characters in the key below.

Key to the Adult *Sandalus* of Virginia  
(after Staines, 1982)

Sides of prothorax uniformly narrowed from posterior to anterior (Fig. 2), keel weakly developed, especially at basal third (Fig. 3) (17-25 mm).....  
.....*Sandalus niger* Knoch

Sides of prothorax subangulate behind middle (Fig. 4), and distinctly keeled throughout (Fig. 5) (12-18 mm).....*Sandalus petrophya* Knoch

*Sandalus niger* Knoch

This species is known from southern Ontario to Florida, west to Colorado and Texas. **Canada:** Ontario (Hicks, 1942). **United States:** Alabama, Colorado, District of Columbia, Florida, Georgia, Indiana, Illinois, Iowa, Kansas, Kentucky, Maryland, Michigan, Missouri, Nebraska, New Jersey, North Carolina, Ohio, Oklahoma, Pennsylvania, South Carolina, South Dakota, Tennessee, Texas, Virginia, West Virginia, and Wisconsin (Brimley, 1938; Downey & Arnett, 1996; Dury, 1879, Elzinga, 1977, Craighead, 1921, Young & Katovich, 2002; Kirk, 1969, 1970; Leng, 1928; Manee, 1908; Rings, 1942; Skelley, pers. comm.; Staines, 1982; Ulke, 1902; Young, 2002). Specimens have been examined from the following counties and cities in **Virginia:** Alexandria (City), Chesapeake (City), Fairfax, Franklin, Montgomery, Richmond (City), Scott, and Shenandoah. (AVEC, GWMP, NMNH, VMNH, VPIC). In Virginia, individuals of *S. niger* have been found from July through November. This species is widespread in Virginia (Fig. 6) and has been collected in the Coastal Plain, Piedmont, Appalachian Plateau, and Valley and Ridge physiographic regions.

*Sandalus petrophya* Knoch  
**NEW STATE RECORD**

This species occurs from New York to Florida, west to Indiana, Missouri, and Alabama, but has not been reported from Virginia previously. **United States:** Alabama, District of Columbia, Florida, Georgia, Illinois, Indiana, Kentucky, Louisiana, Maryland, Missouri, New Jersey, New York, North Carolina, Ohio, Pennsylvania, Tennessee, Virginia, and West Virginia (Blatchley, 1910; Carlton, pers. comm.; Downie & Arnett, 1996; Dury, 1882; Leng, 1928; Say, 1835; Peck & Thomas, 1998; Smith, 1900; Staines, 1982; Ulke, 1902).



Fig. 2. Dorsal view of prothorax of *Sandalus niger* Knoch, male (top) and female (bottom).



Fig. 4. Dorsal view of prothorax of *Sandalus petrophya* Knoch, male (top) and female (bottom).



Fig. 3. Lateral prothoracic margin of *Sandalus niger* Knoch.



Fig. 5. Lateral prothoracic margin of *Sandalus petrophya* Knoch.




Fig. 6. County and city distribution map for *Sandalus niger* Knoch based on specimens examined from Virginia.



Fig. 7. County and city distribution map for *Sandalus petrophya* Knoch based on specimens examined from Virginia.

*Sandalus petrophya* has been found in the following counties and cities in **Virginia**: **Alexandria (City)**: Del Ray, 27 October 2000, R. B. & A. Faden (1); Washington Golf Club, Jewell, 16 August 1925, L.L. Buchanan (1). **Craig Co.**: Jefferson National Forest, FR 620, 12 August 1997, M. W. Donahue (1). **Dickenson Co.**: Breaks Interstate Park, 1-14 July 2000, R. Vigneault (1). **Fairfax Co.**: parking lot, Turkey Run Park Headquarters, McLean, 20 July 2006, B. W. Steury (1); same data, except 28 July 2009 (1); 2 mi. SW Vienna, 30 July 1932, J. C. Bridwell (1); Vienna, 23 July 1913, R. A. Cushman (1); W Falls Church, VI-28-1917, on oak tree, J. L. Wrenn (1); Falls Church, 18 July 1914, H. B. Kirk; same data except 13 August 1914, G. M. Greene (1); same data except 14 August 1914 (2); same data except 7 August 1914 (1); same data except 26 August 1917 (1); Springfield, 26-29 July 1968, Joseph W. Adams (2); Black Pond, 29 June 1914, E. Shoemaker (1); Dead Run, 6 August 1915, R. C. Shannon (1); Great Falls, VII.4 [no year] (1). **Floyd Co.**: county only, 23 August 1982, M. W. Allen (2). **Nelson Co.**: county only, 1 August 1917, W. Robinson (14). **Prince William Co.**: Bull Run Mountains Natural Area Preserve, Little Ridge Loop Trail at Catlett Crk., N38.82642° W077.70203°, 28 July 2008, A.V. Evans, flying/woods AM (1); Occoquan, VIII.1919, Wickham (1). **Rockingham Co.**: Rader Mountain, FR 597, 7 October 1997, M. Donahue (1). **Spotsylvania Co.**: Fredericksburg, 11 August 1897, E. A. Schwarz (1). **Virginia Beach (City)**: Camp Pendleton Annex, vic. NW jct. S. Birdneck & Washington Rds., N36.81383° W075.97079°, Malaise trap, 22 July-19 August 2009, A. V. Evans (1). **Wise Co.**: Powell Mountain Karst Preserve, Cedar Ridge Malaise trap, ca. 1.3 km E Cracker Neck Church, N36.8538533° W082.6998265°, 29 July-20 August 2009, C. S. Hobson (1); same data except 20 August-21 September 2009 (1). (AVEC, GWMP, NMNH, VMNH, VPIC).

Little has been published on this species. Say (1835) frequently observed this species on the flowers of “a

resinous plant common on the prairies of Missouri.” Wenzel (1886) observed *S. petrophya* near Philadelphia during the summer, mostly in mid-July. Dirt encrusted adults, mostly females, were found emerging from the ground in the morning and crawling up the trunks of American Beech (*Fagus grandiflora*). They have also been found under bark or on trunks of trees (Blatchley, 1910).

In Virginia, *S. petrophya* is widespread (Fig. 7) and found in the Coastal Plain, Piedmont, Appalachian Plateau, and Ridge and Valley physiographic regions. Habitat at the Turkey Run Park collection site is successional Tulip-tree (*Liriodendron tulipifera*) forest with nearby mature stands of mesic mixed hardwood forest consisting of oaks (*Quercus* spp.), hickory (*Carya* spp.) and Sugar Maple (*Acer saccharum*). Evans observed and collected a single female flying next to a stream through a wooded area about 1000 h in July in the Bull Run Mountains Natural Area Preserve in Prince William County.

#### ACKNOWLEDGMENTS

We thank Christopher Carlton (Louisiana State Arthropod Museum, Louisiana State University, Baton Rouge, LA), Eric Day (Virginia Polytechnic Institute and State University, Blacksburg, VA), Phil Harpootlian (Simpsonville, SC), Richard Hoffman (Virginia Museum of Natural History, Martinsville, VA), Floyd Shockley (National Museum of Natural History, Washington, DC), Jan Ciegler (Columbus, SC), and Paul Skelley (Florida State Collection of Arthropods, Florida Department of Agriculture-DPI, Gainesville, FL) for supplying collection data or providing access to specimens of *Sandalus* in their care. John Lawrence (Gympie, Queensland, Australia) promptly provided literature pertinent to this study. Fieldwork at the Bull Run Mountain Natural Area Preserve, Camp Pendleton Annex, and Powell Mountain Karst Preserve that produced specimens of

*Sandalus* was conducted by Evans while he was employed as a field zoologist with the Virginia Department of Conservation, Division of Natural Heritage, Richmond, VA. Michael Kieffer (Bull Run Mountains Conservancy, The Plains, Virginia), the Cave Conservancy of the Virginias (Glen Allen, Virginia), and Taura Huxley (NAVFAC Atlantic, Norfolk, VA) provided access and additional logistical support for these surveys. Chris Hobson (Virginia Department of Conservation and Recreation, Division of Natural Heritage, Richmond, VA) selected and established the Malaise trap site in the Powell Mountain Karst Preserve that produced specimens of *Sandalus petrophya*. The National Park Service funded Evans' work on a beetle survey of the George Washington Memorial Parkway during 2010-2011. We also thank Anne Chazal (Virginia Department of Conservation and Recreation, Division of Natural Heritage, Richmond, VA) for generating the distribution maps, Kerry Katovich (University of Wisconsin-Whitewater, Whitewater, WI), Steve Roble (Virginia Department of Conservation and Recreation, Division of Natural Heritage, Richmond, VA), and Charles Staines (National Museum of Natural History, Washington, DC) suggested numerous corrections and comments that improved the accuracy and clarity of this paper.

## LITERATURE CITED

- Arnett, R. H., Jr. 1963. The Beetles of the United States. The Catholic University of America Press, Washington, DC. 1112 pp.
- Blatchley, W. S. 1910. An illustrated catalogue of the Coleoptera or beetles (exclusive of the Rhynchophora) known to occur in Indiana with bibliography and descriptions of new species. Indiana Department of Geology and Natural Resources Bulletin 1: 1-1386.
- Brimley, C. C. 1938. The Insects of North Carolina. Being a List of the Insects of North Carolina and Their Close Relatives. North Carolina Department of Agriculture. 560 pp.
- Craighead, F. C. 1921. Larvae of the North American beetle *Sandalus niger* Knoch. Proceedings of the Entomological Society of Washington 23:44-48.
- Dodge, H. R. 1941. Observations on *Sandalus niger* Knoch, its egg, and first instar larva. Annals of the Entomological Society of America 34: 458-466.
- Downie, N. M., & R. H. Arnett, Jr. 1996. The Beetles of Northeastern North America Volume I. The Sandhill Crane Press, Gainesville, FL. 880 pp.
- Dury, C. 1879. List of the Coleoptera observed in the vicinity of Cincinnati. Journal of the Cincinnati Society of Natural History 2: 162-178.
- Dury, C. 1882. Coleoptera of the vicinity of Cincinnati. Journal of the Cincinnati Society of Natural History 5: 218-220.
- Elzinga, R. J. 1977. Observations on *Sandalus niger* Knoch (Coleoptera: Sandalidae) with description of the triungulin larva. Journal of the Kansas Entomological Society 50: 324-328.
- Hicks, S. D. 1942. Observations on the adult of *Sandalus niger* Knoch in southern Ontario (Coleoptera: Rhipiceridae). Canadian Entomologist 74: 78-79.
- Katovich, K. 2002. 39. Rhipiceridae Latreille, 1834, Pp. 92-94 In R. H. Arnett, Jr., M. C. Thomas, P. E. Skelley, & J. H. Frank (eds.), American Beetles Volume 2, Polyphaga: Scarabaeoidea through Curculionoidea. CRC Press, Washington, DC.
- Kirk, V. M. 1969. A list of the beetles of South Carolina. Part 1-Northern Coastal Plain. Technical Bulletin 1033, South Carolina Agricultural Experimental Station, Clemson, SC. 124 pp.
- Kirk, V. M. 1970. A list of the beetles of South Carolina. Part 2-Mountains, Piedmont, and Southern Coastal plain. Technical Bulletin 1038, South Carolina Agricultural Experimental Station, Clemson, SC. 117 pp.
- Knoch, A. W. 1801. Neue Beyträge zur Insektenkunde. Theil. 1. Schwickertschen Verlage. Leipzig. 208 pp.
- Lawrence, J. F. 2005. 16.2. Rhipiceridae Latreille, 1834. Pp. 456-460 In R. G. Beutel, & R. A. B. Leschen (volume eds.), Coleoptera, Beetles. Volume 1: Morphology and Systematics (Archostemata, Adephaga, Myxophaga, Polyphaga partim). In: N. P. Kristensen, & R. G. Beutel (eds.), Handbook of Zoology. A Natural History of the Phyla of the Animal Kingdom. Volume IV. Arthropoda: Insecta. Part 38. Walter de Gruyter, Berlin and New York.
- LeConte, J. L. 1862. Classification of the Coleoptera of North America. Part 1. Smithsonian Miscellaneous Collections. Washington, DC. 242 pp.
- Leng, C. W. 1928. Order Coleoptera. Pp. 203-520 In

- M. D. Leonard, (ed.), A list of the insects of New York with a list of the spiders and certain other allied groups. Cornell University Agricultural Experiment Station Memoir 101. 1,121 pp.
- Manee, A. H. 1908. Some observations at Southern Pines, N. Carolina. *Entomological News* 19: 286-289.
- Peck, S. B., & M. C. Thomas, 1998. A distributional checklist of the beetles (Coleoptera) of Florida. *Arthropods of Florida and Neighboring Land Areas*. Volume 16. Florida Department of Agriculture and Consumer Services, Gainesville, FL. 180 pp.
- Rings, R. W. 1942. The external anatomy of *Sandalus niger* Knoch (Coleoptera: Rhipiceridae). *Annals of the Entomological Society of America* 35: 411-425.
- Say, T. 1835. Art. X. Descriptions of new North American coleopterous insects, and observations on some already described. *Boston Journal of Natural History* 1: 151-203.
- Smith, J. B. 1900 [1899]. *Insects of New Jersey*. A list of species occurring in New Jersey with notes on those of economic importance. Twenty-seventh Annual Report of the State Board of Agriculture, Supplement. 755 pp.
- Staines, C. L. 1982. The Rhipiceridae of Maryland. *Maryland Entomologist* 2: 38-40.
- Ulke, H. 1902. A list of the beetles of the District of Columbia. *Proceedings of the United States National Museum* 25: 1-57.
- Wenzel, H. W. 1886. Note on *Sandalus*. *Entomological Americana* 1: 107.
- Young, D. K., & K. Katovich. 2002. First record of Rhipiceridae (Coleoptera: Polyphaga: Dascilloidea) from Wisconsin. *Great Lakes Entomologist* 35: 33-34.